arXiv:0907.0251v1 [astro-ph.HE] 1 Jul 2009

A simple model for spectral swapping of supernova neutrinos

Huaiyu Duan

Institute for Nuclear Theory, University of Washington, Seattle, WA 98195, USA

Abstract. Neutrinos emitted from a core-collapse supernova can experience collective flavor transformation because of high neutrino fluxes. As a result, neutrinos of different flavors can have their energy spectra (partially) swapped, a phenomenon known as the (stepwise) spectral swapping or spectral split. We give a brief review of a simple model that explains this phenomenon.

Keywords: supernovae, neutrino oscillations
PACS: 14.60.Pq, 97.60Bw

INTRODUCTION

During a core-collapse supernova event, the neutrino fluxes near the nascent neutron star can be so large that the change of neutrino refractive indices due to neutrino-neutrino forward scattering or neutrino self-interaction becomes important [1, 2]. It was demonstrated in several numerical simulations (e.g., [3, 4, 5, 6]) that neutrinos of different flavors can swap (parts of) their energy spectra as they traverse the supernova envelope (see Figure 1). Here we briefly reviewed a simple model that explains this spectral swap/split phenomenon. (See [7] for a more detailed review.)

With neutrino self-interaction neutrinos with various energies and propagating in different directions can be coupled together, which makes it difficult to analyze neutrino flavor transformation in supernovae. For simplicity we adopt a spherically symmetric supernova model under the “single-angle approximation”. The idea described in the rest of the talk can be generalized to anisotropic environments [8]. In this simple model we assume that the flavor evolution of neutrinos is independent of neutrino propagation directions. The flavor state $|\psi_{\nu,E}(r)\rangle$ of a neutrino ν with energy E at radius r can be solved from the Schrödinger-like equation

$$\mathrm{i}\frac{\mathrm{d}}{\mathrm{d}r}|\psi_{\nu,E}(r)\rangle = \hat{H}|\psi_{\nu,E}(r)\rangle. \tag{1}$$

In the flavor basis the Hamilton in equation (1) can be written as [9, 10]

$$\mathsf{H} = \frac{\mathsf{M}^2}{2E} + \sqrt{2}G_\mathrm{F}\,\mathrm{diag}[n_e(r),0,0] + \sqrt{2}G_\mathrm{F}\int_0^\infty \mathrm{d}E'\,[\rho_{E'}(r) - \bar{\rho}_{E'}(r)], \tag{2}$$

where M is the neutrino mass matrix, G_F is the Fermi’s constant, $n_e(r)$ is the electron number density, and $\rho_{E'}(r)$ and $\bar{\rho}_{E'}(r)$ are the flavor density matrices for the neutrino and the antineutrino with energy E', respectively [11].



FIGURE 1. The initial and final energy spectra of ν_e and ν_x (left panel) and the corresponding (schematic) neutrino survival probability $P_{\nu\nu}$ (right panel) in one of the numerical simulations (single-angle approximation, two-flavor mixing and inverted neutrino mass hierarchy) in [3].

NEUTRINO FLAVOR ISOSPIN

We shall use the two-flavor mixing scheme (ν_e, ν_x) where ν_x is an appropriate mixture of ν_μ and ν_τ. We take the effective vacuum mixing angle and the mass-squared difference to be $\theta_\mathrm{v} \approx \theta_{13} \ll 1$ and $\delta m^2 \approx \pm\delta m^2_\mathrm{atm}$, respectively, where the plus (minus) sign is for the normal (inverted) neutrino mass hierarchy. The two-flavor neutrino oscillation can be visualized with the help of, e.g., the neutrino flavor isospin (NFIS) [12]

$$\vec{s}_\omega = \begin{cases} \psi^\dagger_{\nu,E}(\vec{\sigma}/2)\psi_{\nu,E}, & \text{for neutrinos } (\omega = \delta m^2/2E), \\ (\sigma_y\psi_{\bar\nu,E})^\dagger(\vec{\sigma}/2)(\sigma_y\psi_{\bar\nu,E}), & \text{for antineutrinos } (\omega = -\delta m^2/2E). \end{cases} \tag{3}$$

Equations (1) and (2) now become

$$\frac{\mathrm{d}}{\mathrm{d}r}\vec{s}_\omega = \vec{s}_\omega \times \vec{H}_\omega = \vec{s}_\omega \times (\omega\vec{e}^{\,\mathrm{v}}_z - \sqrt{2}G_\mathrm{F}n_e\vec{e}^{\,\mathrm{f}}_z - \mu\langle\vec{s}\rangle). \tag{4}$$

In equation (4) $\vec{e}^{\,\mathrm{v}}_z$ and $\vec{e}^{\,\mathrm{f}}_z$ are the unit vectors in flavor space along which NFIS's representing $|\nu_1\rangle$ and $|\nu_e\rangle$ must be oriented, respectively. The misalignment of $\vec{e}^{\,\mathrm{v}}_z$ and $\vec{e}^{\,\mathrm{f}}_z$ (with $2\theta_\mathrm{v}$ between them) signifies the non-coincidence of the mass basis and the flavor basis of neutrinos. In vacuum an NFIS $\vec{s}_\omega$ initially aligned with $\vec{e}^{\,\mathrm{f}}_z$ (i.e., a pure ν_e) precesses about $\vec{e}^{\,\mathrm{v}}_z$ with (angular) frequency ω. When projected onto the $\vec{e}^{\,\mathrm{f}}_z$-axis, this precession yields vacuum oscillations with $|\langle\nu_e|\psi_{\nu,E}\rangle|^2 = \frac{1}{2} + \vec{s}_\omega \cdot \vec{e}^{\,\mathrm{f}}_z$. We emphasize that NFIS $\vec{s}_\omega$ precesses in *flavor space* not coordinate space and has nothing to do with the internal spin of the neutrino. Also in equation (4) $\mu = 2\sqrt{2}G_\mathrm{F}n^\mathrm{tot}_\nu$ is the coupling strength between NFIS's, and $\langle\vec{s}\rangle = \int_{-\infty}^{\infty}\mathrm{d}\omega'\, f_{\omega'}\vec{s}_{\omega'}$ is the average NFIS, where n^tot_ν is the total neutrino number density and $f_{\omega'}$ is the (constant) normalized distribution function determined by neutrino energy spectra at the neutrino sphere.

The effects of neutrino masses, ordinary matter and background neutrinos on collective neutrino oscillations can be easily gleaned from equation (4): (a) Because the mass term [the first term of $\vec{H}_\omega$ in equation (4)] is different for neutrinos/antineutrinos with different energies, it induces different precession of the corresponding NFIS's. In other words, the mass term tends to disrupt the collective motion of NFIS's or the collective neutrino oscillation. (b) Because the matter term (the second term of $\vec{H}_\omega$) is independent of ω, it is neutral to collective neutrino oscillations and can be "ignored".[1] (c) The last term of $\vec{H}_\omega$, representing neutrino self-interaction, couples NFIS's with different values of ω and, therefore, drives collective neutrino oscillations.

COLLECTIVE PRECESSION MODE

For simplicity we ignore the matter effect for the reason mentioned above. The equations of motion (4) for NFIS's possess a symmetry about $\vec{e}_z$. (From now on we do not distinguish between $\vec{e}_z^{\,\mathrm{f}}$ and $\vec{e}_z^{\,\mathrm{v}}$ because $\theta_\mathrm{v} \ll 1$.) If a group of NFIS's $\{\vec{s}_\omega | \omega \in (-\infty, +\infty)\}$ satisfies equation (4), then $\{\vec{\tilde{s}}_\omega | \omega \in (-\infty, +\infty)\}$ also obey equation (4), where $\vec{\tilde{s}}_\omega$ are obtained by rotating $\vec{s}_\omega$ about $\vec{e}_z$ by an ω-independent angle ϕ. This symmetry suggests the existence of a collective neutrino oscillation mode in which all the NFIS's precess about $\vec{e}_z$ with the same angular frequency ω_pr [16, 17, 8]. In this collective precession mode and with constant n_ν^tot all NFIS's must be static in the reference frame which rotates about $\vec{e}_z$ with ω_pr. It follows that $\vec{s}_\omega$ must be either aligned or antialigned with $\vec{\tilde{H}}_\omega = \vec{H}_\omega - \omega_\mathrm{pr}\vec{e}_z$, the effective field experience by $\vec{s}_\omega$ in this corotating frame [3], and

$$\vec{s}_\omega = \frac{\epsilon_\omega}{2}\frac{\vec{\tilde{H}}_\omega}{|\vec{\tilde{H}}_\omega|}, \quad \epsilon_\omega = \pm 1. \tag{5}$$

One can then show that all NFIS's in a collective precession mode is fully determined by $\langle s_z \rangle$ and $\langle s_\perp \rangle$, the components of $\langle \vec{s} \rangle$ that are parallel and orthogonal to $\vec{e}_z$, respectively, and the common precession frequency ω_pr. The value of $\langle s_z \rangle$ is fixed by the conservation law that arises from the symmetry described above [13], and ω_pr and $\langle s_\perp \rangle$ can be solved from [18]

$$1 = -\frac{1}{2}\int_{-\infty}^{\infty} \mathrm{d}\omega \frac{\epsilon_\omega f_\omega}{\sqrt{[(\omega - \omega_\mathrm{pr})/\mu - \langle s_z \rangle]^2 + \langle s_\perp \rangle^2}}, \tag{6}$$

$$\omega_\mathrm{pr} = -\frac{1}{2}\int_{-\infty}^{\infty} \mathrm{d}\omega \frac{\omega \epsilon_\omega f_\omega}{\sqrt{[(\omega - \omega_\mathrm{pr})/\mu - \langle s_z \rangle]^2 + \langle s_\perp \rangle^2}}. \tag{7}$$

If n_ν^tot decreases slowly with r, neutrinos will stay in the collective precession mode, and $\vec{s}_\omega$ remains either aligned or antialigned with $\vec{\tilde{H}}_\omega$. Note that this is similar to the adiabatic Mikheyev-Smirnov-Wolfenstein (MSW) flavor transformation [19, 20]. In the adiabatic MSW transformation a neutrino stays in either of the two eigenstates of H_ω (with non-vanishing n_e but zero neutrino flux) which are the two NFIS states aligned and antialigned with $\vec{H}_\omega$, respectively. If the collective precession mode of neutrino oscillations does not break down until $n_\nu^{\rm tot}$ becomes negligible, all the NFIS's will become either aligned or antialigned with $\vec{e}_z$, and

---

[1] The effect of ordinary matter is subtle. When n_e is large, the mass basis is essentially replaced by the flavor basis with a small effective mixing angle no matter what the original value of θ_v is [12, 13, 14]. A large matter density can also suppress collective neutrino oscillations in anisotropic environments [15, 8].

$$\vec{s}_\omega \cdot \vec{e}_z = \frac{\epsilon_\omega}{2}\,\mathrm{sgn}(\omega-\omega^0_{\rm pr}) = \begin{cases} |\langle\nu_e|\psi_{\nu,E}\rangle|^2 - \frac{1}{2} & \text{for neutrino,}\\ \frac{1}{2} - |\langle\bar{\nu}_e|\psi_{\bar{\nu},E}\rangle|^2 & \text{for antineutrino,}\end{cases} \tag{8}$$

where $\omega^0_{\rm pr} = \omega_{\rm pr}|_{n_\nu^{\rm tot}\to 0}$. Correspondingly, $P_{\nu\nu}$, the probability for the neutrino/antineutrino to be in the same flavor as it is at the neutrino sphere, is a step function of ω (the right panel of Figure 1). This is exactly what is observed in numerical simulations (left panel of Figure 1). This phenomenon is called "stepwise spectral swapping" because ν_e and ν_x swap energy spectra at energies above (below) the critical energy $E_{\rm C} = |\delta m^2/2\omega^0_{\rm pr}|$ in an inverted (normal) mass hierarchy case [3]. This is also known as the "spectral split" because $E_{\rm C}$ "splits the transformed spectrum sharply into parts of almost pure but different flavors" [18].

ACKNOWLEDGMENTS

The author would like to thank the conference organizers for the opportunity to give this talk. The author also thank J. Carlson, G. M. Fuller and Y.-Z. Qian for useful discussions. This work is supported in part by US DOE Grant No. DE-FG02-00ER41132 at INT.

REFERENCES

1. G. M. Fuller, R. W. Mayle, J. R. Wilson, and D. N. Schramm, *Astrophys. J.* **322**, 795 (1987).
2. D. Nötzold, and G. Raffelt, *Nucl. Phys.* **B307**, 924 (1988).
3. H. Duan, G. M. Fuller, J. Carlson, and Y.-Z. Qian, *Phys. Rev.* **D74**, 105014 (2006).
4. H. Duan, G. M. Fuller, J. Carlson, and Y.-Z. Qian, *Phys. Rev. Lett.* **97**, 241101 (2006).
5. G. L. Fogli, E. Lisi, A. Marrone, and A. Mirizzi, *JCAP* **0712**, 010 (2007).
6. B. Dasgupta, and A. Dighe, *Phys. Rev.* **D77**, 113002 (2008).
7. H. Duan, and J. P. Kneller (2009), `arXiv:0904.0974`.
8. H. Duan, G. M. Fuller, and Y.-Z. Qian (2008), `arXiv:0808.2046`.
9. Y. Z. Qian, and G. M. Fuller, *Phys. Rev.* **D51**, 1479 (1995).
10. J. T. Pantaleone, *Phys. Rev.* **D46**, 510 (1992).
11. G. Sigl, and G. Raffelt, *Nucl. Phys.* **B406**, 423 (1993).
12. H. Duan, G. M. Fuller, and Y.-Z. Qian, *Phys. Rev.* **D74**, 123004 (2006).
13. S. Hannestad, G. G. Raffelt, G. Sigl, and Y. Y. Y. Wong, *Phys. Rev.* **D74**, 105010 (2006).
14. H. Duan, G. M. Fuller, and Y.-Z. Qian, *Phys. Rev.* **D77**, 085016 (2008).
15. A. Esteban-Pretel, et al., *Phys. Rev.* **D78**, 085012 (2008).
16. H. Duan, G. M. Fuller, J. Carlson, and Y.-Z. Qian, *Phys. Rev.* **D75**, 125005 (2007).
17. H. Duan, G. M. Fuller, and Y.-Z. Qian, *Phys. Rev.* **D76**, 085013 (2007).
18. G. G. Raffelt, and A. Y. Smirnov, *Phys. Rev.* **D76**, 081301(R) (2007).
19. L. Wolfenstein, *Phys. Rev.* **D17**, 2369 (1978).
20. S. P. Mikheyev, and A. Y. Smirnov, *Yad. Fiz.* **42**, 1441 (1985), [Sov. J. Nucl. Phys. 42, 913 (1985)].